arXiv:1401.7511v1 [math.CO] 29 Jan 2014

FURTHER INEQUALITIES BETWEEN VERTEX-DEGREE-BASED TOPOLOGICAL INDICES

AKBAR ALI, AKHLAQ AHMAD BHATTI AND ZAHID RAZA

Department of Mathematics
National University of Computer and Emerging Sciences
Block-B, Faisal Town, Lahore, Pakistan
akbarali.maths@gmail.com, akhlaq.ahmad@nu.edu.pk, zahid.raza@nu.edu.pk

Abstract. Continuing the recent work of L. Zhong and K. Xu [*MATCH Commun. Math. Comput. Chem.* **71** (2014) 627-642], we determine inequalities among several vertex-degree-based topological indices; first geometric-arithmetic index *(GA)*, augmented Zagreb index *(AZI)*, Randić index *(R)*, atom-bond connectivity index *(ABC)*, sum-connectivity index *(X)* and harmonic index *(H)*.

1. Introduction

Let $G = (V, E)$ denote a simple graph with vertex set $V(G) = \{v_1, v_2, ..., v_n\}$ and edge set $E(G)$ such that $|E(G)| = m$. Suppose that d_i is the degree of a vertex $v_i \in V(G)$ and ij is edge connecting the vertices v_i and v_j [1].

Topological indices are numerical parameters of a graph which are invariant under graph isomorphisms. They play a significant role in mathematical chemistry especially in the QSPR/QSAR investigations [2, 3].

L. Zhong and K. Xu [29] obtained several inequalities among R, ABC, X and H indices. An important topological index that was not discussed in [29] is the AZI index. B. Furtula et al.[30] proved that AZI index is a valuable predictive index in the study of the heat

Table 1. Degree-based topological indices discussed in this paper

Name of index	Definition of index
Randić*(R)*, [4]-[8]	$R(G) = \sum_{ij \in E(G)} \frac{1}{\sqrt{d_i d_j}}$
Harmonic*(H)*, [9]-[12]	$H(G) = \sum_{ij \in E(G)} \frac{2}{d_i + d_j}$
Atom-bond connectivity*(ABC)*, [13]-[19]	$ABC(G) = \sum_{ij \in E(G)} \sqrt{\frac{d_i + d_j - 2}{d_i d_j}}$
Sum-connectivity*(X)*, [20]-[26]	$X(G) = \sum_{ij \in E(G)} \frac{1}{\sqrt{d_i + d_j}}$
First geometric-arithmetic*(GA)*, [27, 28]	$GA(G) = \sum_{ij \in E(G)} \frac{\sqrt{d_i d_j}}{\frac{1}{2}(d_i + d_j)}$
Augmented Zagreb *(AZI)*, [30]-[33]	$AZI(G) = \sum_{ij \in E(G)} \left(\frac{d_i d_j}{d_i + d_j - 2} \right)^3$

of formation in octanes and heptanes. I. Gutman and J. Tošovič [33] recently tested the correlation abilities of 20 vertex-degree-based topological indices for the case of standard heats of formation and normal boiling points of octane isomers, and they found that the augmented Zagreb index yield the best results. GA index is another important topological index, not discussed in [29]. It has been demonstrated, on the example of octane isomers, that GA index is well-correlated with a variety of physico-chemical properties [27]. For the mathematical properties of the GA index and their applications in QSPR and QSAR see the survey[28] and the references cited therein. In this note, we continue the work of L. Zhong and K. Xu [29] and establish some inequalities among the topological indices given in Table 1.

2. Inequalities Between Vertex-Degree-Based Topological Indices

In this section, we give inequalities among several vertex-degree-based topological indices such as AZI, GA, R, ABC, X and H indices.

Theorem 2.1. *If G is a connected graph with $n \geq 2$ vertices, then*

$$\sqrt{2}X(G) \leq GA(G) \leq \sqrt{2(n-1)}X(G).$$

The lower bound is attained if and only if $G \cong P_2$ and the upper bound is attained if and only if $G \cong K_n$.

Proof. Without loss of generality we can assume $1 \leq d_i \leq d_j \leq n-1$. Consider the function

$$F(x,y) = \left(\frac{\frac{2\sqrt{xy}}{x+y}}{\frac{1}{\sqrt{x+y}}}\right)^2 = \frac{4xy}{x+y} \quad \text{where} \quad 1 \leq x \leq y \leq n-1$$

One can easily see that $F(x,y)$ is strictly monotone increasing in both x and y. This implies that $F(x,y)$ attains the minimum value at $(x,y) = (1,1)$ and the maximum value at $(x,y) = (n-1, n-1)$. Hence

$$2 = F(1,1) \leq F(x,y) \leq F(n-1,n-1) = 2(n-1)$$

which implies,

$$\sqrt{2} \leq \frac{GA(G)}{X(G)} \leq \sqrt{2(n-1)}$$

with the left equality if and only if $(d_i, d_j) = (1,1)$ for every edge ij of G and the right equality if and only if $(d_i, d_j) = (n-1, n-1)$ for every edge ij of G. This completes the proof. □

If graph G has the minimum degree at least 2, then the lower bound in Theorem 2.1 can be improved:

Corollary 2.2. *If G is a connected graph with minimum degree $\delta \geq 2$, then*

$$\sqrt{2\delta}X(G) \leq GA(G)$$

with equality if and only if G is a δ-regular graph.

Theorem 2.3. *If G is a connected graph with $n \geq 2$ vertices, then*

$$R(G) \leq GA(G) \leq (n-1)R(G).$$

The lower bound is attained if and only if $G \cong P_2$ and the upper bound is attained if and only if $G \cong K_n$.

Proof. Suppose that $1 \leq d_i \leq d_j \leq n-1$ and let

$$F(x,y) = \frac{\frac{2\sqrt{xy}}{x+y}}{\frac{1}{\sqrt{xy}}} = \frac{2xy}{x+y} \quad \text{where} \quad 1 \leq x \leq y \leq n-1.$$

It can be easily seen that $F(x,y)$ is strictly monotone increasing in both x and y. This implies that

$$1 = F(1,1) \leq F(x,y) \leq F(n-1,n-1) = n-1$$

therefore,

$$1 \leq \frac{GA(G)}{R(G)} \leq (n-1)$$

with the left equality if and only if $(d_i, d_j) = (1,1)$ for every edge ij of G and the right equality if and only if $(d_i, d_j) = (n-1, n-1)$ for every edge ij of G. □

If the graph G has the minimum degree $\delta \geq 2$, then the lower bound in Theorem 2.3 can be replaced by $\delta R(G)$.

Corollary 2.4. *If G is a connected graph with $\delta \geq 2$, then*

$$\delta R(G) \leq GA(G)$$

with equality if and only if G is a δ-regular graph.

B. Zhou and N. Trinajstić [23] proved that if G is a connected graph with $n \geq 3$ vertices, then $\sqrt{\frac{2}{3}}R(G) \leq X(G)$ with equality if and only if $G \cong P_3$. Hence from Theorem 2.1, we have:

Corollary 2.5. *If G is a connected graph with with $n \geq 3$ vertices, then*

$$\sqrt{\frac{4}{3}}R(G) \leq GA(G) \leq (n-1)R(G)$$

with equality if and only if $G \cong P_3$ and right equality if and only if $G \cong K_n$.

Theorem 2.6. *If G is a connected graph with $n \geq 2$ vertices, then*

$$H(G) \leq GA(G) \leq (n-1)H(G).$$

The lower bound is attained if and only if $G \cong P_2$ and the upper bound is attained if and only if $G \cong K_n$.

Proof. Using the same technique, used in proving Theorem 2.1 and Theorem 2.3, one can easily prove the required result. □

Corollary 2.7. *If G is a connected graph with minimum degree $\delta \geq 2$, then*

$$\delta H(G) \leq GA(G) \tag{2.1}$$

with equality if and only if G is a δ-regular graph

Theorem 2.8. *If G is a connected graph having $n \geq 3$ vertices with minimum degree $\delta \geq 2$, then*

$$\frac{\sqrt{2(n-2)}}{n-1}GA(G) \leq ABC(G) \leq \frac{n+1}{4\sqrt{n-1}}GA(G) \tag{2.2}$$

with left equality if and only if $G \cong K_n$ and right equality if and only if $G \cong C_3$.

Proof. Suppose that $2 \leq d_j \leq d_i \leq n-1$ and consider the function

$$F(x,y) = \left(\frac{\sqrt{\frac{x+y-2}{xy}}}{\frac{2\sqrt{xy}}{x+y}} \right)^2 = \frac{(x+y)^2(x+y-2)}{4x^2y^2} \quad \text{where} \quad 2 \leq y \leq x \leq n-1.$$

Then

$$\frac{\partial F(x,y)}{\partial y} = -\frac{(x+y)\{x^2-y^2+x(x+y-4)\}}{4x^2y^3} \leq 0$$

This implies that $F(x,y)$ is monotone decreasing in y. Hence $F(x,y)$ attains the maximum value at $(x,y) = (x,2)$ for some $2 \leq x \leq n-1$.
But,

$$\frac{dF(x,2)}{dx} = \frac{x(x^2-4)}{16x^3} \geq 0$$

that is, $F(x,2)$ is monotone increasing in x which implies $F(x,2)$ has maximum value at $x = n-1$. Hence

$$F(x,y) \leq F(n-1,2) = \frac{(n+1)^2}{16(n-1)}$$

and therefore,

$$\frac{ABC(G)}{GA(G)} \leq \frac{n+1}{4\sqrt{n-1}}$$

with the equality if and only if $(d_i, d_j) = (n-1, 2)$ for every edge ij of G, i.e.,

$$ABC(G) \leq \frac{n+1}{4\sqrt{n-1}} GA(G)$$

with the equality if and only if $G \cong C_3$
Since $F(x,y)$ is monotonously decreasing in y. It means $F(x,y)$ attains minimum value at (x,x) for some $2 \leq x \leq n-1$. Since

$$\frac{dF(x,x)}{dx} = \frac{2x(2-x)}{x^4} \leq 0$$

hence,

$$\frac{2(n-2)}{(n-1)^2} = F(n-1,n-1) \leq F(x,x) \leq F(x,y)$$

i.e.,

$$\frac{\sqrt{2(n-2)}}{(n-1)} \leq \frac{ABC(G)}{GA(G)}$$

with the equality if and only if $(d_i, d_j) = (n-1, n-1)$ for every edge ij of G, which completes the proof. □

L. Zhong and K. Xu [29] proved that if $\delta \geq 2$ in a connected graph G, then

$$H(G) \leq R(G) \leq X(G) < ABC(G) \tag{2.3}$$

with the first equality if and only if G is a regular graph, and the second equality if and only if G is a cycle. Hence, from Theorem 2.8 and inequality (2.3), we have:

Corollary 2.9. *If G is a connected graph with minimum degree $\delta \geq 2$, then*

$$H(G) \leq R(G) \leq X(G) < ABC(G) \leq \frac{n+1}{4\sqrt{n-1}} GA(G)$$

with the first equality if and only if G is a regular graph, the second equality if and only if G is a cycle, and last with equality if and only if $G \cong C_3$

Denoted by T^* the tree on eight vertices, obtained by joining the central vertices of two copies of star $K_{1,3}$ by an edge. K. C. Das and N. Trinajstić [18] proved that

$$GA(G) > ABC(G) \tag{2.4}$$

for every molecular graph $G \ncong K_{1,4}, T^*$. The same authors proved that inequality (2.4) holds for any graph $G \ncong K_{1,4}, T^*$ in which $\Delta - \delta \leq 3$. In [19], it is proved that if $\delta \geq 2$ and $\Delta - \delta \leq (2\delta - 1)^2$ then inequality (2.4) holds.

Corollary 2.10. *If G is a connected graph satisfying at least one of the following properties:*

(i): *G is molecular graph such that $G \ncong K_{1,4}, T^*$*
(ii): *$\Delta - \delta \leq 3$ and $G \ncong K_{1,4}, T^*$*
(iii): *$\delta \geq 2$ and $\Delta - \delta \leq (2\delta - 1)^2$,*

then

$$H(G) \leq R(G) \leq X(G) < ABC(G) < GA(G)$$

Denote the chromatic number of a graph G by $\chi(G)$. Deng et al. [11] proved that for every connected graph G

$$\chi(G) \leq 2H(G) \tag{2.5}$$

with equality if and only if G is a complete graph. From inequalities (2.1) and (2.5), we obtain a sharp upper bound of $\chi(G)$ in terms of GA index:

Corollary 2.11. *If G is a connected graph of order n with minimum degree $\delta \geq 2$, then*

$$\chi(G) \leq \frac{2}{\delta} GA(G)$$

with equality if and only if $G \cong K_n$.

Another vertex-degree-based topological Index is the modified second Zagreb index defined [34, 35] as:

$$M_2^*(G) = \sum_{ij \in E(G)} \frac{1}{d_i d_j}$$

Using the same technique, used in proving Theorem 2.1 and Theorem 2.3, one can easily prove the following result:

Theorem 2.12. *If G is a connected graph with $n \geq 2$ vertices, then*

$$M_2^*(G) \leq R(G) \leq (n-1) M_2^*(G), \tag{2.6}$$

$$\frac{M_2^*(G)}{\sqrt{2}} \leq X(G) \leq \frac{(n-1)^{\frac{3}{2}}}{\sqrt{2}} M_2^*(G), \tag{2.7}$$

$$M_2^*(G) \leq H(G) \leq (n-1) M_2^*(G), \tag{2.8}$$

$$M_2^*(G) \leq GA(G) \leq (n-1)^2 M_2^*(G), \tag{2.9}$$

$$\sqrt{2} M_2^*(G) \leq ABC(G) \leq (n-1)\sqrt{2(n-2)} M_2^*(G); n \geq 3. \tag{2.10}$$

The left equality in (2.6)-(2.9) and in (2.10) is attained if and only if $G \cong P_2$ and $G \cong P_3$ respectively. The right equality in all inequalities (2.6)-(2.10) is attained if and only if $G \cong K_n$.

Corollary 2.13. *If G is a connected graph with minimum degree $\delta \geq 2$, then*

$$\delta M_2^*(G) \leq R(G), \tag{2.11}$$

$$\frac{\delta^{\frac{3}{2}} M_2^*(G)}{\sqrt{2}} \leq X(G), \tag{2.12}$$

$$\sqrt{\delta} M_2^*(G) \leq H(G), \tag{2.13}$$

$$\delta^2 M_2^*(G) \leq GA(G), \tag{2.14}$$

$$\delta \sqrt{2(\delta-1)} M_2^*(G) \leq ABC(G). \tag{2.15}$$

The equality in all inequalities (2.11)-(2.15) is attained if and only if G is a δ-regular graph.

Now, we establish some inequalities between augmented Zagreb index and other vertex-degree-based topological indices.

Theorem 2.14. *If G is a connected graph having $n \geq 3$ vertices, then*

$$\frac{1536}{343} X(G) \leq AZI(G) \leq \frac{\sqrt{(n-1)^{13}}}{\sqrt{32}(n-2)^3} X(G)$$

with left equality if and only if $G \cong S_{1,8}$ and right equality if and only if $G \cong K_n$.

Proof. Without loss of generality we can assume $1 \leq d_i \leq d_j \leq n-1$. Consider the function

$$F(x,y) = \left(\frac{\left(\frac{xy}{x+y-2}\right)^3}{\frac{1}{\sqrt{x+y}}} \right)^2 = (x+y)\left(\frac{xy}{x+y-2}\right)^6 \quad \text{with} \quad 1 \leq x \leq y \leq n-1 \quad \text{and} \quad y \geq 2.$$

Then,

$$\frac{\partial F(x,y)}{\partial x} = \frac{x^5 y^6 \{x^2 + (y-2)(6y+7x)\}}{(x+y-2)^7} \geq 0.$$

This means $F(x,y)$ is increasing in x and hence is minimum at $(1,y_1)$ and maximum at (y_2,y_2) for some $2 \leq y_1, y_2 \leq n-1$. Now,

$$\frac{dF(1,y)}{dy} = \frac{y^5(y^2-7y-6)}{(y-1)^7}$$

and this implies, $F(1,y)$ is monotone decreasing in $2 \leq y \leq 7$ and monotone increasing in $8 \leq y \leq n-1$. Hence minimum value of $F(x,y)$ is

$$\min\{F(1,7), F(1,8)\} = 9\left(\frac{8}{7}\right)^6$$

Therefore,

$$3\left(\frac{8}{7}\right)^3 \leq \frac{AZI(G)}{X(G)} \tag{2.16}$$

with equality if and only if $(d_i, d_j) = (1,8)$ for each edge ij of G.
Moreover,

$$F(y,y) = \frac{y^{13}}{32(y-1)^6}$$

is monotone increasing and hence

$$F(x,y) \leq F(n-1,n-1) = \frac{(n-1)^{13}}{32(n-2)^6}$$

therefore,

$$\frac{AZI(G)}{X(G)} \leq \frac{\sqrt{(n-1)^{13}}}{\sqrt{32}(n-2)^3} \tag{2.17}$$

with equality if and only if $(d_i, d_j) = (n-1, n-1)$ for each edge ij of G. From (2.16) and (2.17), required result follows. □

If graph G has the minimum degree at least 2, then the lower bound in Theorem 2.14 can be improved:

Corollary 2.15. *If G is a connected graph with minimum degree $\delta \geq 2$, then*

$$\frac{\delta^{\frac{13}{2}}}{\sqrt{32}(\delta-1)^3} X(G) \leq AZI(G)$$

with equality if and only if G is a δ-regular graph.

Using the similar technique, used in proving Theorem 2.14, one can prove the following result (we omit the proof)

Theorem 2.16. *If G is a connected graph with $n \geq 3$ vertices, then*

$$\frac{343\sqrt{7}}{216} R(G) \leq AZI(G) \leq \frac{(n-1)^7}{8(n-2)^3} R(G), \tag{2.18}$$

$$\frac{375}{64} H(G) \leq AZI(G) \leq \frac{(n-1)^7}{8(n-2)^3} H(G), \tag{2.19}$$

$$\left(\frac{n-1}{n-2}\right)^{\frac{7}{2}} ABC(G) \leq AZI(G) \leq \left(\frac{(n-1)^2}{2(n-2)}\right)^{\frac{7}{2}} ABC(G), \tag{2.20}$$

$$8GA(G) \leq AZI(G) \leq \frac{(n-1)^6}{8(n-2)^3} GA(G); \quad \delta \geq 2, \tag{2.21}$$

$$4M_2^*(G) \leq AZI(G) \leq \frac{(n-1)^4}{2(n-2)} M_2^*(G). \tag{2.22}$$

The left equality in (2.18), (2.19), (2.20), (2.21), (2.22) hold if and only if $G \cong S_{1,7}$, $G \cong S_{1,5}$, $G \cong S_{1,n-1}$, $G \cong C_n$, $G \cong P_3$ respectively and right equality if and only if $G \cong K_n$.

Corollary 2.17. *If G is a connected graph with $\delta \geq 2$, then*

$$\frac{\delta^7}{8(\delta-1)^3} R(G) \leq AZI(G), \tag{2.23}$$

$$\frac{\delta^7}{8(\delta-1)^3} H(G) \leq AZI(G), \tag{2.24}$$

$$\left(\frac{\delta^2}{2(\delta-1)}\right)^{\frac{7}{2}} ABC(G) \leq AZI(G), \tag{2.25}$$

$$\frac{\delta^6}{8(\delta-1)^3}GA(G) \leq AZI(G), \tag{2.26}$$

$$\frac{\delta^4}{2(\delta-1)}M_2^*(G) \leq AZI(G). \tag{2.27}$$

The equality in all inequalities (2.23)-(2.27) hold if and only if G is a δ-regular graph.

References

[1] F. Harary, *Graph Theory*, Addison-Wesley, Reading, MA, 1969.
[2] J. Devillers, A.T. Balaban (Eds.), *Topological Indices and Related Descriptors in QSAR and QSPR*, Gordon and Breach, Amsterdam, 1999.
[3] I. Gutman, B. Furtula (Eds.), *Novel Molecular Structure DescriptorsTheory and Applications vols. III*, Univ. Kragujevac, Kragujevac, 2010.
[4] M. Randić, On characterization of molecular branching, *J. Am. Chem. Soc.* **97** (1975) 6609-6615.
[5] L. B. Kier, L. H. Hall, *Molecular Connectivity in Structure-Activity Analysis*, Wiley, New York, 1986.
[6] L. Pogliani, From molecular connectivity indices to semiempirical connectivity terms: Recent trends in graph theoretical descriptors, *Chem. Rev.* **100** (2000) 3827-3858.
[7] X. Li and Y. Shi, A survey on the Randic index, *MATCH Commun. Math. Comput. Chem.* **59** (2008), 127-156.
[8] B. Liu et al., On the conjecture of Aouchiche and Hansen about the Randic index, *Discrete Math.* **313** (2013) 225-235
[9] S. Fajtlowicz, On conjectures of Grati-II, *Congr. Numer.* **60** (1987) 187-197.
[10] R. Wua, Z. Tanga, H. Deng, A lower bound for the harmonic index of a graph with minimum degree at least two, *Filomat* **27** (2013), 51-55
[11] H. Deng, S. Balachandran, S. K. Ayyaswamy, Y. B. Venkatakrishnan, On the harmonic index and the chromatic number of a graph, *Discrete Appl. Math.* **161** (2013) 2740-2744.
[12] L. Zhong, The harmonic index for graphs, *Appl. Math. Lett.* **25** (2012) 561-566.
[13] E. Estrada, L. Torres, L. Rodrguez, I. Gutman, An atom-bond connectivity index: modelling the enthalpy of formation of alkanes, *Indian J. Chem. A* **37** (1998) 849-855.
[14] E. Estrada, Atom-bond connectivity and the energetic of branched alkanes, *Chem. Phys. Lett.* **463** (2008) 422-425.
[15] S. A. Hosseini, M. B. Ahmadi, I. Gutman, Kragujevac trees with minimal atom-bond connectivity index, *MATCH Commun. Math. Comput. Chem.* **71** (2014) 5-20.
[16] L. Gan, H. Hou, B. Liu, Some results on atom-bond connectivity index of graphs, *MATCH Commun. Math. Comput. Chem.* **66** (2011) 669-680.
[17] W. Lin, T. Gao, Q. Chen, X. Lin, On the minimal ABC index of connected graphs with given degree sequence, *MATCH Commun. Math. Comput. Chem.* **69** (2013) 571-578.
[18] K. C. Das, N. Trinajstić, Comparison between first geometric-arithmetic index and atom-bond connectivity index, *Chem. Phys. Lett.* **497** (2010) 149-151.
[19] Z. Raza, A.A. Bhatti, A. Ali, On "comparison between first geometric-arithmetic index and atom-bond connectivity Index", submitted.
[20] B. Zhou, N. Trinajstić, On a novel connectivity index, *J. Math. Chem.* **46** (2009) 1252-1270.
[21] B. Lučić, S. Nikolić, N. Trinajstić, B. Zhou, S. Ivaniš Turk, Sum-connectivity index, in: I. Gutman, B. Furtula (Eds.), *Novel Molecular Structure Descriptors-Theory and Applications I*, Univ. Kragujevac, Kragujevac, 2010, pp. 101-136.
[22] B. Lučić, Trinajstić, B. Zhou, Comparison between the sum-connectivity index and product-connectivity index for benzenoid hydrocarbons, *Chem. Phys. Lett.* **475** (2009) 146-148.
[23] B. Zhou, N. Trinajstić, Relations between the product and sum-connectivity indices, *Croat. Chem. Acta* **85** (2012) 363-365.
[24] R. Xing, B. Zhou, N. Trinajstić, Sum-connectivity index of molecular trees, *J. Math. Chem.* **48** (2010) 583-591.
[25] S. Wang, B. Zhou, N. Trinajstić, On the sum-connectivity index, *Filomat* **25** (2011) 29-42.
[26] Z. Du, B. Zhou, Trinajstić, Minimum sum-connectivity indices of trees and unicyclic graphs of a given matching number, *J. Math. Chem.* **47** (2010) 842-855.

[27] D. Vukičević , B. Furtula, Topological index based on the ratios of geometrical and arithmetical means of end-vertex degrees of edges, *J. Math. Chem.* **46** (2009) 1369-1376.
[28] K. C. Das, I. Gutman, and B. Furtula, Survey on geometric-arithmetic indices of graphs, *MATCH Commun. Math. Comput. Chem.* **65** (2011) 595-644.
[29] L. Zhong and K. Xu, Inequalities between vertex-degree-based topological Indices, *MATCH Commun. Math. Comput. Chem.* **71** (2014) 627-642
[30] B. Furtula, A. Graovac, D. Vukicevic, Augmented Zagreb index, *J. Math. Chem.* **48** (2010) 370-380
[31] Y. Huang, B. Liu, L. Gan, Augmented zagreb index of connected graphs, *MATCH Commun. Math. Comput. Chem.* **67** (2012) 483-494
[32] D. Wang, Y. Huang, B. Liu, Bounds on augmented zagreb index, *MATCH Commun. Math. Comput. Chem.* **68** (2012) 209-216
[33] I. Gutman, J. Tošovič, Testing the quality of molecular structure descriptors: Vertex-degree-based topological indices, *J. Serb. Chem. Soc.* **78** (2013) 805-810
[34] S. Nikolić, G. Kovačević, A. Miličević, N. Trinajstić, The zagreb indices 30 years after, *Croat. Chem. Acta* **76** (2003) 113-124
[35] A. Miličević, S. Nikolić, On variable Zagreb indices, *Croat. Chem. Acta* **77** (2004) 97-101.